

Assignments \mathcal{N}^o 3

released: 12.11.2014 **due:** 18.11.2014 at 12:00h

Task 1: Dependency in ERGM

5 points

Let \mathcal{G} be the set of undirected, loopless graphs with $n = 3$ vertices and consider an exponential random graph model (\mathcal{G}, P) with only one statistic, namely $t(G)$ (the number of triangles) with associated parameter value $\log(3)$.

Compute the edge probability of a dyad and the conditional edge probability, given that there is an edge on another dyad. Are dyads independent or not in this model?

Task 2: Inappropriate Sampling from ERGM

5 points

Provide a (preferably simple) example of an ERGM (\mathcal{G}, P) which demonstrates that the following algorithm returns graphs G with probability **different from** $P(G)$.

Algorithm 1: incorrect ERGM sampling

Data : edge set E

Output: random graph $G = (\{1, \dots, n\}, E)$

$E \leftarrow \emptyset$

foreach $e \in D = \{d_1, \dots, d_{\binom{n}{2}}\}$ **do**

 randomly add e to E with probability

$$\frac{P(V, E \cup \{e\})}{P(V, E) + P(V, E \cup \{e\})}$$

Hints: use a small model in which you have dependency among dyads. Show that the algorithm turns the very first dyad d_1 into an edge with a probability that is different from the probability $P(d_1 \in E)$ defined by the ERGM. You might reuse probability-calculations from the lecture or from another task.

Task 3: Estimating ERGM parameters

10 points

Download the *Preprocessed Knecht Classroom Data* from <http://www.inf.uni-konstanz.de/algo/lehre/ws14/nm/local/data/data.html>.

- (a) Import the adjacency matrix of the network observed at the third time point (file `net-3.csv`) and the demographic characteristics of the actors (file `demographics.csv`)
 - (a.1) Symmetrize the adjacency matrix using the function `symmetrize` in the R package `sna`. Use the “strong” rule.
 - (a.2) Create a network object using the symmetrized matrix and name the object `netwundir`. Check that the network is undirected and add the gender of the pupils as an attribute.
 - (a.3) Estimate an ERGM model specified only by the number of edges and interpret the result.
 - (a.4) Estimate an ERGM model specified by the number of: edges, triangles, 2-stars, 3-stars and the same gender covariate. Interpret the result.
- (b) The data are directed. During the lecture, several statistics were introduced for undirected ties. Analogous statistics exist for directed ties. Therefore, we can estimate ERGMs also for directed data.
 - (b.1) Create a network object using the adjacency matrix observed at the third time point. Name the object `netwdir`. Check that the network is directed and add the gender of the pupils as an attribute.
 - (b.2) Estimate an ERGM specified by the following statistics: number of edges, reciprocal dyads, and homophilous dyads with respect to gender. Interpret the result.
 - (b.3) Estimate an ERGM specified by the following statistics: number of edges, reciprocal dyads, homophilous dyads with respect to

gender and transitive triplets. What can you observe from the result?

Hint: estimation of this model won't converge.

N.b.: To find the name of the effects check the help `?ergm` and the link `ergm-terms`.